

Virbac

ECTOPARASITICIDE

FOR EXTERNAL ANIMAL USE ONLY

Amidip® Max

Reg. No. G3767 (Act 36/1947)

EMULSIFIABLE CONCENTRATE

Controls ticks, kills lice and mange mites on cattle.

Contra-indications: Do not use on horses.

OXPECKER COMPATIBLE

CAUTION

■ STORAGE INSTRUCTIONS

Store below 25 °C. Protect product from extremes of heat and cold and from dampness.

■ COMPOSITION

Amitraz 25,0 % m/v

■ WARNINGS

- Withdrawal period: do not slaughter animals for human consumption within 7 days of last treatment.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this label available to him.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

- Ensure that the container remains tightly sealed.
- Destroy empty containers.
- Wash hands thoroughly after use.
-

■ DIRECTIONS FOR USE:

Use only as directed.

For external use only.

Spray race: Once prepared, the spray tank solution should be used within a period of 3 days.

AMIDIP MAX should be incorporated into the spray race according to the following dilution:

AMIDIP MAX	WATER
100 ml	100 L
200 ml	200 L
500 ml	500 L
1 L	1 000 L
5 L	5 000 L
10 L	10 000 L
20 L	20 000 L

Dilute the required quantity of AMIDIP MAX in approximately 10 L water, and then incorporate the AMIDIP MAX solution into the required volume of water in the spray race. Mix thoroughly.

Boost: It is recommended that not more than 500 head be sprayed in the spray race at one time, particularly where fouling is a problem. Where this figure is exceeded, boosting at the rate of 100 ml per 100 head commencing at 500 head can be undertaken to maintain a satisfactory deposit of active ingredient, especially where tick-borne diseases are a problem.

Plunge dip: Once prepared, the dip tank solution should be used within a period of 7 days.

AMIDIP MAX should be incorporated into the dip tank at a rate of 1 L AMIDIP MAX per 5 000 L water.

Total replenishment: The dip tank should be made up to volume prior to each dipping interval. AMIDIP MAX should be added with each dipping interval at a rate of 1 L to 5 000 L, taking the total volume of water in the tank into consideration.

Treatment frequency:

Ticks and mange: Spray / dip weekly
Lice: Spray / dip twice with a 7-day interval

In-use combination of AMIDIP MAX (G3767) and Pro-Dip Cyp 20 % (G2311)

CATTLE DIP

HARMFUL

Composition:

AMIDIP MAX:

Amitraz 25 % m/v

Pro-Dip Cyp 20 % (G2311):

Cypermethrin 20 % m/v

■ **INDICATIONS:** The combination of AMIDIP MAX (G3767) and Pro-Dip Cyp 20 % (G2311) is highly effective in the control of ticks.

■ ADDITIONAL WARNINGS:

- Do not slaughter animals for human consumption within 5 days of last treatment.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this label available.

■ ADDITIONAL PRECAUTIONS:

While the safety and stability of an in-use combination of AMIDIP MAX and Pro-Dip Cyp 20% has been established, AMIDIP MAX should not be mixed with any other dip.

■ PRESENTATION

100 ml, 1 L, and 5 L fluorinated HDPE bottles.

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000

Namibia Reg. No. V04/18.3.4/131 NS0
Botswana Reg. No. W130660