

**EFFIPRO SPOT-ON FOR SMALL DOGS**

Reg. No. G4052 (Act 36/1947)

EFFIPRO SPOT-ON FOR MEDIUM DOGS

Reg. No. G4056 (Act 36/1947)

EFFIPRO SPOT-ON FOR LARGE DOGS

Reg. No. G4051 (Act 36/1947)

EFFIPRO SPOT-ON FOR EXTRA LARGE DOGS

Reg. No. G4057 (Act 36/1947)

CAUTION**■ STORAGE INSTRUCTIONS**

Store below 30°C in a dry place

■ COMPOSITION

Each ml contains 100 mg fipronil

■ INDICATIONS

For the treatment and prevention of infestations by fleas (*Ctenocephalides felis*) and the tick, (*Rhipicephalus sanguineus*) in dogs. Use as an aid in the treatment of flea allergy dermatitis (FAD) in dogs, where this has been previously diagnosed by a veterinarian.

■ WARNINGS

Do not use on puppies less than 2 months old and/or weighing less than 2 kg. Do not use on sick or convalescing animals. Do not use in rabbits, as adverse reactions and even death could occur. This product is specifically developed for dogs. Do not use in cats, as this could lead to overdosing. Keep out of reach of children, uninformed persons and animals. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

This product can cause mucous membrane and eye irritation. Contact of the product with mouth and eyes should be avoided. In the case of accidental eye contact, immediately and thoroughly flush the eyes with water. If eye irritation persists, seek medical advice and show the package insert to the physician or veterinarian. Do not smoke, drink or eat during application. Avoid contact with fingers. If this occurs, wash hands thoroughly with soap and water. Always wash hands after use. Animals or operators with a known hypersensitivity to insecticides or alcohol should avoid contact with the product. Do not overdose. Should accidental ingestion occur, contact a doctor or veterinarian immediately. It is important to make sure that the product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment. If licking occurs, a brief period of hypersalivation may be observed, due mainly to the nature of the carrier. Treated animals should not be handled until the application site is dry. It is recommended that animals be treated in the early evening and that recently treated animals not be allowed to sleep or play with owners,

especially children. The product contains alcohol; therefore, keep treated animals away from fires and other sources of ignition for at least 30 minutes or until the hair coat is dry. The alcohol carrier may have adverse effects on painted, varnished or other household surfaces or furnishings. Dispose of empty pipettes with domestic refuse.

■ DOSAGE AND DIRECTIONS FOR USE - Use only as directed

Animals should be weighed accurately prior to treatment.

Administer by topical application to the skin.

Dose: 1 pipette of 0,67 ml per dog, weighing over 2 kg and up to 10 kg. 1 pipette of 1,34 ml per dog weighing over 10 kg and up to 20 kg. 1 pipette of 2,68 ml per dog weighing over 20 kg and up to 40 kg. 1 pipette of 4,02 ml per dog weighing over 40 kg and up to 60 kg

Method of administration: Hold the pipette upright. Tap the narrow part of the pipette to ensure that the contents are within the main body of the pipette. Break the snap-off top of the spot-on pipette along the scored line. Part the dog's coat until the skin is visible. Place the tip of the pipette directly against the bare skin and squeeze gently several times to empty its contents. Repeat this procedure at one or two different points along the dog's back.

Avoid bathing or shampooing the dog one hour before, or 2 days after, application of EFFIPRO, as this may affect the efficacy of the product. After the dog has been treated with EFFIPRO, do not bath or shampoo the dog more than once a week. The minimum treatment interval is 4 weeks. When used as part of a strategy for the treatment of flea allergy dermatitis, monthly applications to the allergic patient and to other dogs in the household are recommended. For optimal control of flea infestations in multi-pet households, all dogs in the household should be treated with a suitable insecticide.

Residual effect: EFFIPRO is effective against new flea infestations for approximately 3 months. The product has a persistent acaricidal efficacy of up to 4 weeks against ticks. If ticks of the species *Rhipicephalus sanguineus* are present when the product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week.

■ PRESENTATION Cartons of 1 x 0,67 / 1,34 / 2,68 or 4,02 ml pipettes

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115, Halfway House 1685,
South Africa
Tel.: +27 12 657-6000