

FOR ANIMAL USE ONLY

②+⑧†

Prazifen® Tablets

Reg. No. G2602 (Act 36/1947)

Treatment for roundworm, hookworm, whipworm and tapeworm infestation in dogs and cats**■ STORAGE INSTRUCTIONS**

Store at room temperature below 25 °C

■ COMPOSITION

Each tablet contains:

Praziquantel 50 mg
Fenbendazole 500 mg**■ WARNINGS**

- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

In cases of overdosage, reactions of discomfort, vomiting and diarrhoea may occur. If these symptoms are observed, consult your veterinarian. Treatment is symptomatic and supportive.

■ EFFICACY

For the treatment of mixed infections with roundworm, hookworm and tapeworm caused by:

Roundworms - *Toxocara canis*
 Toxascaris leonina
Hookworms - *Ancylostoma caninum*
Whipworm - *Trichuris vulpis*
Tapeworms - *Echinococcus* spp.
 Dipylidium caninum
 Taenia spp.

■ DIRECTIONS FOR USE

Use only as directed

Tablets to be administered orally. The recommended dosage is 5 mg praziquantel and 50 mg fenbendazole per kg body mass, i.e. one tablet per 10 kg body mass.

To ease administration, tablets can be mixed with a small portion of food. Dietary measures or fasting are not necessary. To ease dispensing to clients, the tablets are scored; give a quarter of tablet per 2,5 kg body mass.

■ PRESENTATION

Blister pack of 12 tablets.
Cartons containing 96 tablets.

†Benzimidazoles + isoquinolones

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115, Halfway House 1685,
South Africa
Tel: +27 12 657 6000

Namibia Reg. No. V02/18.1.8/20 NSO