



## ENDOPARASITICIDE

FOR EXTERNAL ANIMAL USE ONLY

# Flukazole C

Reg. No. G3533 (Act 36/1947)

FOR THE CONTROL OF LIVER FLUKE, ROUNDWORMS, LUNG- WORMS AND TAPEWORMS IN CATTLE AND SHEEP.

### ■ STORAGE INSTRUCTIONS

Store below 30 °C in a cool dry place.

### CAUTION

#### ■ COMPOSITION

Triclabendazole ..... 12 % m/v  
Oxfendazole ..... 4,53 % m/v

#### ■ WARNINGS

- Do not slaughter animals for human consumption within 28 days of dosing.
- Do not dose lactating cows whose milk or milk products are intended for human consumption.
- Do not use less than 21 days before calving/lambing, in animals where milk and milk products from treated animals may be used for human consumption.
- Where accidental treatment with FLUKAZOLE C takes place within the 21-day period, milk must not be used for human consumption, or be supplied for processing for at least 21 days following treatment. Young animals fed this milk within 21 days of slaughter must not be used for human consumption.
- Containers and any residual contents should be disposed of safely in accordance with national waste disposal regulations. Empty containers should not be reused for any other purpose.
- Dangerous to fish. DO NOT contaminate dams, rivers or streams with the drench or used container.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

#### ■ PRECAUTIONS

- Wash hands after use.
- Avoid contact with eyes and skin. May cause irritation.
- Rinse the medicated drench three times after use. Do not dispose of undiluted chemicals.
- Wash equipment thoroughly after use.
- **First aid:** If swallowed, and if more than 15 minutes from a hospital, induce vomiting.

#### ■ DIRECTIONS FOR USE

Use only as directed Shake before use.

**Cattle and Sheep:** Dosage: 1 ml / 10 kg body mass orally. This dose delivers 12 mg Triclabendazole per kg body mass and 4,53 mg Oxfendazole per kg body mass. Cattle in excess of 650 kg body mass should be dosed at 5 ml per 50 kg body mass. Dose the group according to the body mass of the heaviest animal in the group.

#### Efficacy:

**Roundworms:** FLUKAZOLE C controls benzimidazole-sensitive mature and immature roundworms and lungworms.

Roundworm Species- Cattle	Immature	Adult
Wireworm ( <i>Haemonchus placei</i> )	*	*
Brown Stomachworm ( <i>Ostertagia ostertagi</i> )	*	*
Cattle Bankruptworm ( <i>Cooperia</i> spp.)	*	*
Hookworm ( <i>Bunostomum phlebotomum</i> )	*	*
Nodular worm ( <i>Oesophagostomum radiatum</i> )	*	*
Lungworm ( <i>Dictyocaulus viviparous</i> )	*	*

Definition: \* Controls (≥ 90 % effective)

Roundworm Species- Sheep	Immature	Adult
Wireworm ( <i>Haemonchus placei</i> )	*	*
Brown Stomachworm ( <i>Ostertagia ostertagi</i> )	*	*
Large mouth bowel worm ( <i>Chabertia</i> spp)	*	*
Long-necked bankrupt worm ( <i>Nematodirus</i> spp)	*	*
Lung worm ( <i>Dictyocaulus</i> spp)	*	*
Bankrupt worm ( <i>Trichostrongylus</i> spp)	*	*
Hookworm ( <i>Gaigeria</i> spp)	*	*
White bankrupt worm ( <i>Strongyloides</i> spp)	*	*

Definition: \* Controls (≥ 90 % effective)

**Ovicidal:** Roundworm eggs present in the animal at the time of dosing are prevented from hatching. Cattle and sheep should not be placed on clean pasture until 6 hours after drenching.

**Tapeworm:** FLUKAZOLE C also controls tapeworm in cattle and sheep.

**Milk tapworm** (*Moniezia* spp).

Class 1 (100 % effective in 80% of the treated animals).

**Liver Fluke:** FLUKAZOLE C is used for the treatment of susceptible early immature, immature and mature Liver Fluke.

FLUKAZOLE C controls the following:

	Early immatures		Late immatures			Adult
	2 weeks	3 weeks	4 weeks	6 weeks	8 weeks	
Liver Fluke ( <i>Fasciola hepatica</i> )	*		*	*		*
Giant Liver Fluke ( <i>Fasciola gigantica</i> )		*		*		*

Definition: \* Controls (≥ 90 % effective)

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information contact your veterinarian.

#### ■ PRESENTATION

250 ml, 1 L and 5 L opaque HDPE bottles with polypropylene cap.

#### ■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd  
(Reg. No. 1990/003743/07)  
Private Bag X115  
Halfway House 1685  
South Africa  
Tel.: +27 12 657 6000

Namibia Reg. No. V06/18.1.8/76 NS0  
Zambia MA No: 359/713V **P-V**