

FOR ANIMAL USE ONLY

Virbacox

Reg. No. G2266 (Act 36/1947)

An aid in the prevention of coccidiosis in broiler chickens caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. burnetti* and *E. mivati*.

POISONOUS

■ STORAGE INSTRUCTIONS

Store in a cool dry place. Close container after use.

■ COMPOSITION

Salinomycin sodium 12 % m/m

■ WARNINGS

- Do not feed to laying or breeding hens, turkeys, horses or other equines. Ingestion by horses can be fatal.
- Birds must not be slaughtered for human consumption during treatment or for at least 5 days after the last treatment.
- Birds should not be treated with products containing tiamulin for at least 7 days before or after receiving feed containing this product. Severe growth depression or death may result.
- Do not leave or dispose of VIRBACOX, or feed containing it, where it may be accessible to animals not under treatment, or to wildlife.
- Handle with care. Poisonous when swallowed.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ Precautions:

- Operators should avoid direct contact with the product or inhalation.
- Wear face masks to at least the specification of BS 2091, impervious gloves and protective overalls when handling.
- Rinse eyes thoroughly with water immediately if contact with eyes occurs.
- Medicated feed must be used within 3 months of mixing.

■ DIRECTIONS FOR USE

Use only as directed

- Mix 450 g to 550 g VIRBACOX per ton of complete feed to supply 54 to 66 mg / kg Salinomycin sodium.
- Normal dosage rate is 500 g per ton providing 60 mg / kg of Salinomycin sodium.

■ PRESENTATION

Containers of 25 kg

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000