

ANTIPARASITIC

①*

FOR ANIMAL USE ONLY

Virbamax Premix

Reg. No. G3122 (Act 36/1947)

FOR THE PREVENTION AND TREATMENT OF INTERNAL AND EXTERNAL PARASITES IN PIGS.

CAUTION

■ **STORAGE INSTRUCTIONS**

Store at room temperature below 25 °C.

■ **COMPOSITION**

Abamectin 0,6 % m/m

■ **WARNINGS**

- Do not slaughter animals intended for human consumption within 14 days of treatment.
- Pigs of any age, pregnant sows and breeders can be treated with VIRBAMAX PREMIX.
- It is recommended to start the treatment of sows 2 to 3 weeks before farrowing.
- Use in other animals, especially dogs, may cause severe reactions including fatalities.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ **PRECAUTIONS**

- Take adequate measures not to create dust when incorporating the product into feed.
- Wear a dust mask, gloves and approved safety glasses.
- Avoid contact with the skin and eyes. Rinse in case of exposure. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
- Do not smoke, drink or eat when working with the product.
- Dispose of containers and any residual contents safely.

■ **DIRECTIONS FOR USE - Use only as directed**

VIRBAMAX PREMIX should be mixed into the feed of animals being treated. A feed intake of 100 µg abamectin per kg body mass per day is recommended. This is equivalent to 16,7 mg VIRBAMAX PREMIX.

Pigs:

Body mass (kg)	Quantity premix required (g / ton of feed)
< 39	333
40 - 69	400
70 - 100	450

Breeding sows and boars: Include 1,67 kg to 2,50 kg VIRBAMAX PREMIX per ton of feed for consumption at a rate of 1 kg to 1,5 kg per 100 kg body mass daily for 7 consecutive days. The medicated feed can be given ad libitum as the only ration.

Sows:

Feed intake (kg/day)	Body mass (kg)	Quantity premix required (kg/ ton feed)	Number of sows treated per ton of feed for 7 days
2,0	200	1,666	71
	250	2,083	
	300	2,500	
2,5	200	1,333	57
	250	1,667	
	300	2,000	
3	200	1,111	47
	250	1,389	
	300	1,667	

Efficacy: VIRBAMAX PREMIX is effective in the prevention and treatment of common pathogenic infestations due to sensitive strains of the following adult and immature internal and external parasites of swine.

- **Gastro-intestinal roundworms:**
Hyostromylus rubidus, Oesophagostomum spp., Strongyloides ransomi, Stephanurus dentatus and Trichuris suis
- **Ascariidiosis:**
Ascaris suum
- **Lungworm:**
Metastrongyles spp. (adult)
- **Mange:**
Sarcoptes scabiei var suis
- **Lice:**
Hematopinus suis

■ **PRESENTATION**

333 g white, opaque sachet and a 5 kg transparent polyethylene bag in a white polypropylene bucket.

■ **REGISTRATION HOLDER**

VIRBAC RSA (Pty) Ltd
(Reg No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000

* Macrocytic Lactoes