

FOR EXTERNAL ANIMAL USE ONLY

Oberdine

Reg. No. G0495 (Act 36/1947)

Wound, Eye and Footrot Spray.

For the treatment of wounds, cuts, sores, burns, abrasions, shearing wounds and footrot in cattle, sheep, goats, pigs and horses and ringworm in cattle. Also treats infectious ophthalmia and bacterial eye infection in cattle, sheep, goats and horses. Acts in the presence of blood, pus and dead tissue.

COMPOSITION:

Povidone-iodine 10 % m/v Allantoin 2 % m/v Glycerine 5 % m/v

WARNINGS

Keep out of reach of children, uninformed persons and animals. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected seek veterinary advice and notify the registration holder

DIRECTIONS FOR USE – Use only as directed

Wounds, cuts, sores, burns and abrasions: Wash affected area and spray whole area from a distance of 15 – 20 mm.

Dressing: Wash affected area – pour on or apply liberally with brush or cotton wool.

Footrot in cattle, sheep, goats and pigs: Wash affected area and pare off diseased tissue. Spray or pour on. Where possible, cover with dressing. Treat 1 – 2 times daily.

Infectious ophthalmia (Rickettsia) and bacterial eye infections:

Cattle, sheep, goats and horses can be treated by holding the container 15 – 20 mm away from the eye and by spraying gently onto the eye. Repeat treatment twice daily until signs of acute inflammation are controlled, i.e. lacrimation (tear flow), redness and pain have abated. Possible white discolouration of the cornea will gradually disappear by itself.

Ringworm in cattle: Break or scrub off crusts. Apply twice daily for at least 14 days

REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd, (Reg No. 1990/003743/07) Private Bag X115, Halfway House 1685 South Africa,

Tel: +27 12 657 6000