

FOR ANIMAL USE ONLY



Pro-Inject[®] Yellow

Reg. No. G2048 (Act 36/1947)

INJECTABLE SOLUTION

Liver fluke and roundworm remedy for cattle with residual action against re-infestation of wireworm, hookworm and nodular worm

CAUTION

■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C

■ COMPOSITION

Closantel 10 mg/ml

■ WARNINGS

- Allow 30 days between last injection and slaughter for human consumption.
- Not for use in milk cows whose milk is intended for human consumption.
- Handle with care. Poisonous by swallowing and inhalation.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

- Prevent contamination of food, food utensils, etc.
- Wash hands thoroughly after use.
- Use sterile injection apparatus. As for all injections, a sharp, sterile needle should be used for each animal.
- Do not inject animals whilst their coats are wet (e.g. after dipping) due to the risk of infection at the injection site.
- As swelling is known to occur at the site of injection following subcutaneous administration, deep intramuscular administration is preferable.

■ ADMINISTRATION

Administer by subcutaneous injection at side of neck or by intramuscular injection in any large muscle group using a 15-16 G needle 2,5 cm in length. Do not exceed 10 ml per injection site.

DIRECTIONS FOR USE

Use only as directed

1 ml per 40 kg body mass

Dosage table

Body mass (kg)	Dose (ml)
80	2
120	3
180	4,5
200	5
250	6,25
300	7,5

Double dose

When a higher efficacy against internal parasites is indicated, a dose of 1 ml per 20 kg body mass can be administered, e.g.:

Body mass (kg)	Dose (ml)
80	4
120	6
180	9
200	10
250	12,5
300	15

■ EFFICACY

Standard dose at 1 ml per 40 kg body mass

1. Roundworm Species

PRO-INJECT YELLOW controls and aids in the control of the following:

	Im-matures	Adult	Persistence ^{††}
Wireworm	*	*	3 weeks
Hookworm	**	*	-
Nodular Worm	*	*	-

Definition: * Control (≥ 90 % effective)

** Aids in control of (60 - 89 %)

†† Interval between treatment and re-infestation

2. Liver Fluke

PRO-INJECT YELLOW controls the following:

Liver Fluke	4 weeks	6 weeks	Adult
Giant Liver Fluke	6 weeks	8 weeks	Adult

Definition: * Control (≥ 90 % effective)

Double dose at 1 ml per 20 kg body mass

1. Roundworm Species

PRO-INJECT YELLOW controls the following:

	Im-matures	Adult	Persistence ^{††}
Wireworm	*	*	6 weeks
Hookworm	*	*	3 weeks
Nodular Worm	*	*	3 weeks

Definition: * Control (≥ 90 % effective)

†† Interval between treatment and re-infestation

2. Liver Fluke

PRO-INJECT YELLOW controls and aids in the control of the following:

Liver Fluke	4 weeks	6 weeks	Adult
	**	**	*
Giant Liver Fluke	6 weeks	8 weeks	Adult
	**	*	*

Definition: * Control (≥ 90 % effective)

** Aids in control of (60 - 89 %)

■ PRESENTATION

500 ml

† Halogenated salicylanilides

Registration holder:

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115, Halfway House 1685,
South Africa
Tel: +27 12 657 6000