



Prodose[®] Blue

FOR ANIMAL USE ONLY

Prodose[®] Blue

Reg. No. G2090 (Act 36/1947)

②

Oral Suspension

Roundworm and milk tapeworm remedy for cattle

CAUTION

STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

COMPOSITION

Albendazole 7,5 % m/v

WARNINGS:

- Do not slaughter animals for human consumption within 12 days of dosing.
- Do not dose lactating cows whose milk or milk products are intended for human consumption.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS:

- Management practices should ensure that drenching procedures are carefully carried out.
- It is important not to exceed the recommended dosage in cows during the first month of pregnancy.

DIRECTIONS FOR USE:

Use only as directed

Shake well before use

Dosage: The dose is 1 ml per 10 kg body mass, e.g.:

Body mass (kg)	Dose (ml)
100	10
150	15

200	20
250	25
300	30

EFFICACY:

1. Roundworms:

Prodose Blue controls the following:

	Immature	Adult
Wireworm	*	*
Brown Stomachworm	*	*
Cattle Bankruptworm	*	*
Hookworm	*	*
Nodular Worm	*	*

Definition: * Control (≥ 90 % effective)

2. Tapeworms: Milk tapeworm (*Moniezia spp.*)

Class 1 (100 % effective in 80 % of the treated animals)

3. Ovicidal:

Roundworm eggs present in the animal at the time of dosing are prevented from hatching.

[†] Benzimidazoles

PRESENTATION:

200 ml, 1 kg, 5 kg, 10 kg and 20 kg.

REGISTRATION HOLDER:

VIRBAC RSA (Pty) Ltd,
(Reg No. 1990/003743/07)
Private Bag X115,
Halfway House, 1685, South Africa,
Tel: +27 12 657 6000

Namibia No. V95/18.1.1/41 [NSO]

Zimbabwe Reg. No: 2017/80.12.10/9770 (V.M.G.D) 80.10.12 Anthelmintics used in livestock