



# Prodose® Yellow LA

FOR ANIMAL USE ONLY

## Prodose® Yellow LA

Reg. No. G1959 (Act 36/1947)

④

Oral Solution

Liver fluke, conical fluke, nasal worm and roundworm remedy for sheep and goats with residual efficacy against re-infestation of wireworm and hookworm.

CAUTION

### STORAGE INSTRUCTIONS:

Store at room temperature below 25 °C.

### COMPOSITION:

Closantel 7,5 % *m/v*

### WARNINGS:

- Allow 30 days between last dosing and slaughter for human consumption.
- Do not use in milk goats whose milk is intended for human consumption.
- Do not administer to Angora goats before they have been weaned.
- Handle with care.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### PRECAUTIONS:

- Do not repeat treatments more frequently than every 5 weeks.
- As with all worm remedies, it is advisable to ensure animals have been immunised against pulpy kidney disease before drenching.
- Prevent contamination of food, food utensils, etc.
- Wash hands thoroughly after use.

### DIRECTIONS FOR USE - Use only as directed

Dosage table: 1 ml per 10 kg body mass, e.g.

Body mass (kg)	Dosage (ml)
10	1
20	2
30	3
40	4
50	5
60	6

### Immature conical fluke:

Use 1,3 ml per 10 kg body mass as recommended dosage rate.

### EFFICACY:

#### 1. Nasal Worm (*Oestrus ovis*)

Highly effective against 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> stages instar larvae.

#### 2. Roundworms

Prodose Yellow LA Oral Solution controls and aids in the control of the following:

	Immatures	Adults	Persistence <sup>††</sup>
Wireworm	*	*	5 weeks
Hookworm	**	**	2 weeks

Definition: \* Controls (≥ 90 % effective)

\*\* Aids in control of (60 – 89 %)

<sup>††</sup> Interval between treatment and re-infestation

#### 3. Liver Fluke

Prodose Yellow LA Oral Solution controls and aids in the control of the following:

	4 weeks	6 weeks	Adults
Liver fluke		**	*
	6 weeks	8 weeks	Adults
Giant liver fluke		**	*
	Immatures		
Conical Fluke		*	

Definition: \* Control (≥ 90 % effective)

\*\* Aids in control of (60 – 89 %)

<sup>†</sup> Halogenated salicylanilides

### PRESENTATION:

200 ml, 1 kg, 5 kg and 10 kg.

### REGISTRATION NUMBER:

VIRBAC RSA (Pty) Ltd  
 (Reg No. 1990/003743/07)  
 Private Bag X115  
 Halfway House  
 1685  
 South Africa  
 Tel: +27 12 657 6000

Namibia Reg. No: V03/18.1.3/104 NS0