

ENDOPARASITICIDES FOR ANIMAL USE ONLY

③†

Prodose® Red

Reg. No. G2012 (Act 36/1947)

ROUNDWORM REMEDY FOR SHEEP, GOATS, CATTLE AND OSTRICHES

CAUTION

■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

■ COMPOSITION

Levamisole hydrochloride 2,56 % m/v

■ WARNINGS

- Do not slaughter animals for human consumption with-in 7 days of treatment.
- No worm remedy, not even Prodose Red (levamisole) should be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxaemia).
- Although Prodose Red can be administered to pregnant animals, care must be exercised in the handling of animals in the late stage of pregnancy.
- In cattle, the odd animal when treated with levamisole, may develop an allergic type reaction of a transient nature which normally disappears within 24 hours. If necessary an antihistamine may be administered. Under no circumstances must cortisone preparations be used.

- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ DIRECTIONS FOR USE - Use only as directed

Use only as a drench. Dose animals according to the following schedule.

Cattle: 15 ml / 50 kg body mass, for example:

Body mass	Dose
100 kg	30 ml
150 kg	45 ml
200 kg	65 ml
250 kg	75 ml
300 kg	90 ml

Sheep and goats: 3 ml / 10 kg body mass, for example:

Body mass	Dose
10 kg	3 ml
20 kg	6 ml
30 kg	9 ml
40 kg	12 ml

Ostriches: 6 ml / 5 kg body mass, for example:

Body mass	Dose
5 kg	6 ml
50 kg	60 ml
150 kg	180 ml

Safety: Cattle, sheep and goats: Pregnant ewes and cows, lambs, kids and calves may be dosed with Prodose Red.

Ostriches: Important: Hold the bird's head up until the dose has been swallowed. Ensure that the dose is given directly into the gullet and not into the windpipe.

■ EFFICACY

Cattle: Roundworm

	Immature	Adult
Wireworm	*	*
Cattle bankrupt-worm	*	*
Hookworm	-	*
Nodular worm	-	*

Definition: * Controls (≥ 90 % effective)

	Immature	Adult
Brown stomachworm	-	*
Nodular worm	*	*

Definition: * Aids in the control (60-89 % effective)

Sheep and goats: Roundworms

Worm species	Immature	Adult
Wireworm	*	*
Brown stomachworm	-	*
Bankruptworm	*	*
Long-necked bankrupt-worm	*	*
Hookworm	*	*
Nodular worm	*	*
Large-mouthed bowel-worm	*	*
Lungworm	**	**

Definition: * Control (≥ 90% effective)

** Aids in the control (60-89 % effective)

Ostriches: Roundworms

Worm species	Immature	Adult
<i>Libyostrongylus douglassii</i> (Wireworm)	-	*
<i>Codiostormum</i> spp	-	*
<i>Libyostrongylus douglassii</i> (Wireworm)	**	-
<i>Codiostormum</i> spp	**	-

Definition: * Control (≥ 90% effective)

** Aids in the control (60-89 % effective)

■ PRESENTATION

1 L, 5 L polypropylene bottles.

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000

† Imidazoles