

FOR ANIMAL USE ONLY

Replensol

Reg. No. G0550 (Act 36/1947)

An aid in the treatment of all forms of scours in farm animals to prevent dehydration and the loss of essential minerals and energy.

■ STORAGE INSTRUCTIONS

Store in a cool, dry place.

■ COMPOSITION:

Sodium chloride	2,0%
potassium chloride	2,6%
citric acid	3,1%
potassium phosphate	2,3%
glucose	90%
Vitamin A	100 000 IU

■ WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Cattle:

Following poisoning with arsenic, tulip and other irritant plant substances. Prolonged transport by truck or rail usually decreases the water intake, and increases loss of minerals and fluid due to scours caused by stress. Refusal to eat or drink for any reason whatsoever causes fluid and mineral loss which would have been compensated for by daily food and water intake. In many calf diseases accompanied by scours as little time as possible must lapse between onset of symptoms and the administration of Replensol. Sick calves will take Replensol readily while refusing milk and water.

Horses:

Persistent diarrhoea of unknown causes, sometimes lasting many months. Replensol is an ideal form of supportive treatment in these cases and should be given daily in drinking water. As in calves, foals often suffer from scours and as in all young animals treatment with Replensol should

commence immediately and only be stopped two days after recovery.

Pigs:

Many conditions affecting the digestive tract of the pig (particularly young pigs) result in diarrhoea. Even if not fatal, any prolonged case of scours will retard growth to an uneconomical level. Early treatment with Replensol reduces deaths and loss of condition and hastens recovery.

All farm animals, especially the young, are affected by scours at some stage.

■ DIRECTIONS FOR USE

Use only as directed

Adult Cattle and Horses: Dose 15-20 litre through stomach tube or offer instead of drinking water.

Calves and Foals: Dose one litre every 4 hours or offer from bottle with teat.

Pigs (all ages): Offer instead of drinking water.

Adult Sheep: · Dose one litre every 4 hours through stomach tube.

Lambs: Offer 100 - 250 ml every four hours from bottle with teat.

Directions for use: Dissolve 250 g of Replensol in 4,5 litres of lukewarm water. Make available during and for one day after diarrhoea has stopped.

Dosage:

Adult cattle and horses: Dose 15 - 20 litres through stomach tube, or offer instead of drinking water.

Calves and foals: Dose 1 litre every 4 hours, or offer from bottle with teat.

Pigs: Offer instead of drinking water.

Adult sheep: Dose 1 litre every 4 hours through stomach tube.

Lambs: Offer 100 - 250 ml every 4 hours from bottle with teat.

■ PRESENTATION

4 x 250 g

■ REGISTRATION HOLDER

Oberon Pharm (Pty) Ltd
(Reg. No. 1971/009791/07)
P.O. Box 38558
Booysens
2016
South Africa

Namibia Reg. No: N-SR 1592