

## ECTOPARASITICIDES

FOR EXTERNAL ANIMAL USE ONLY

**Stopatik®**

Reg. No. G1431 (Act 36/1947)

**CONTROLS TICKS, HOUSE FLIES, STABLE FLIES AND FACE FLIES IN CATTLE.**

**CONTROLS KAROO PARALYSIS TICK AND BONTLEGGED TICK IN SHEEP.**

**OXPECKER COMPATIBLE**

**CAUTION**

### ■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

### ■ COMPOSITION

Cypermethrin ..... 2 % m/v  
Piperonyl butoxide ..... 8 % m/v

### ■ WARNINGS

- Cattle and sheep must not be slaughtered for human consumption within 28 days of last treatment. There is no withdrawal period of milk for human consumption.
- Handle with care. Poisonous when swallowed.
- Store in a safe place away from food and feed.
- Keep out of reach of children, uninformed persons and pets.
- Toxic to bees, other insects and fish.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek advice and notify veterinary the registration holder.

**NOTE TO DOCTOR:** In case of accidental poisoning or ingestion, consult a medical practitioner and make this information available to him. No specific antidote, treat symptomatically.

### ■ PRECAUTIONS

- Do not mix STOPATIK with any other product.
- Avoid skin and eye contact. Wash hands immediately with soap and water after accidental contact. Contact with the product may cause a burning sensation of the skin and eyes which may last for 24 hours. This reaction is transient and without harmful effect.
- A mild skin sensitivity may occur in certain individual animals.
- Wash hands thoroughly before eating, drinking or smoking.
- Prevent contamination of food, food utensils, drinking water, rivers, dams, grazing, etc.
- Destroy empty containers after use. Never re-use containers for any other purpose.

### ■ DIRECTIONS FOR USE

**Use only as directed.**

**Cattle:** Apply 10 ml / 100 kg body mass to back midline of cattle being treated. Application should be done by means of a suitable applicator from the head to the base of the tail. Treat weekly for three weeks. Then treatment may be extended to once every second week, depending on the fly and tick problem.

**Sheep:** Apply the total dose required in 4 equal portions to both axillae and both sides of the groin. Treat at 4 - weekly intervals during the risk period. Use 1 ml per 10 kg body mass for wool sheep where the wool length is less than 25 mm, and 2 ml per 10 kg mass in hair sheep and wool sheep where the wool length is greater than 25 mm.

### Dosing table: Cattle:

Body mass (kg)	Dose (ml)	Doses per 1 litre	Doses per 20 litre
50	5	200	4 000
100	10	100	2 000
200	20	50	1 000
300	30	33	666
400	40	25	500
500	50	20	400

### Dosing table: Sheep Wool Sheep

Wool length < 25 mm	1 ml per 10 kg body mass
Wool length > 25 mm	2 ml per 10 kg body mass
Hair sheep	2 ml per 10 kg body mass

### ■ PRESENTATION

1L and 20L containers.

### ■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd  
(Reg. No. 1990/003743/07)  
Private Bag X115  
Halfway House  
1685  
South Africa  
Tel: +27 12 657 6000