



Terralon[®] LA

FOR ANIMAL USE ONLY

Terralon[®] LA

Reg. No. G2676 (Act 36/1947)

Injectable Solution

For the treatment of heartwater, tickborne gallsickness (anaplasmosis), pneumonia, footrot, joint-ill, navel-ill and pink eye (infectious bovine keratoconjunctivitis) in cattle, sheep, goats and pigs

CAUTION

STORAGE INSTRUCTIONS:

Store below 25 °C and protect from light

COMPOSITION:

Each ml contains:

Oxytetracycline (as dihydrate) 200 mg

INDICATIONS:

Terralon LA is indicated for the treatment of infections caused by bacteria and micro-organisms susceptible to oxytetracycline in cattle, pigs, sheep and goats.

WARNINGS:

Animals intended for human consumption should not, after last treatment, be slaughtered or the milk used within the time period indicated below

SPECIES	MEAT	MILK
Cattle	30 days	6 days
Pigs	22 days	-
Sheep	22 days	6 days
Goats	30 days	8 days

- Do not use in cats, dogs and horses.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS:

- Contra-indicated in cases of known hypersensitivity, impaired liver and/or kidney function.
- Use only when clearly indicated in the last half of pregnancy.

N.B. Although studies indicate that Terralon LA is well tolerated, a transient irritation may occur at the site of the injection. In lactating animals a small drop in milk production (up to 3 days) may be observed. Hypersensitivity reactions occur occasionally.

Note to veterinarian:

In case of a serious anaphylactic reaction, the administration of epinephrine, antihistamines and corticosteroids should be considered.

DIRECTIONS FOR USE:

Use only as directed

Terralon LA is for deep intramuscular injection only. It is strongly recommended to divide the intramuscular dosages

over two or more injection sites – maximum 15 ml per injection site in cattle over 150 kg, and 7 ml in pigs, calves, sheep and goats. Injection sites should be alternated.

Cattle	1 ml/10 kg body mass (equivalent to 20 mg/kg body mass) If necessary repeat after 72 hours.
Sheep	1 ml/10 kg body mass (equivalent to 20 mg/kg body mass) If necessary repeat after 60 hours.
Goats	1 ml/10 kg body mass (equivalent to 20 mg/kg body mass) If necessary repeat after 72 hours.
Pigs	1 ml/10 kg body mass (equivalent to 20 mg/kg body mass) If necessary repeat after 72 hours.

In case of tickborne gallsickness (anaplasmosis), repetitive treatments (10-14 day treatment period) are necessary to eliminate the parasite from carriers.

PRESENTATION:

Amber coloured glass vials containing 100 ml, 250 ml and 500 ml sterile solution.

REGISTRATION HOLDER:

VIRBAC RSA (Pty) Ltd
(Reg No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: (012) 657 6000

Namibia Reg. No. V15/17.1.2/1297 NSO
Zambia MA No: 359/734V **POM-V**