

FOR ANIMAL USE ONLY

Terralon® LA

Reg. No. G2676 (Act 36/1947)

INJECTABLE SOLUTION

For the treatment of heartwater, tickborne gallsickness (anaplasmosis), pneumonia, footrot, joint-ill, navel-ill and pink eye (infectious bovine keratoconjunctivitis) in cattle, sheep, goats and pigs.

CAUTION**■ STORAGE INSTRUCTIONS**

Store below 25 °C and protect from light

■ COMPOSITION

Each ml contains:

Oxytetracycline (as dihydrate).....200 mg

■ INDICATIONS

TERRALON LA is indicated for the treatment of infections caused by bacteria and micro-organisms susceptible to oxytetracycline in cattle, pigs, sheep and goats.

■ WARNINGS

Animals intended for human consumption should not, after last treatment, be slaughtered or the milk used within the time period indicated below.

SPECIES	MEAT	MILK
Cattle	30 days	6 days
Pigs	22 days	-
Sheep	22 days	6 days
Goats	30 days	8 days

- Do not use in cats, dogs and horses.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS

- Contra-indicated in cases of known hypersensitivity, impaired liver and / or kidney function.
- Use only when clearly indicated in the last half of pregnancy.

N.B. Although studies indicate that TERRALON LA is well tolerated, a transient irritation may occur at the site of the injection. In lactating animals a small drop in milk production (up to 3 days) may be observed. Hypersensitivity reactions occur occasionally.

Note to veterinarian:

In case of a serious anaphylactic reaction, the administration of epinephrine, antihistamines and corticosteroids should be considered.

■ DIRECTIONS FOR USE**Use only as directed**

TERRALON LA is for deep intramuscular injection only. It is strongly recommended to divide the intramuscular dosages over two or more injection sites – maximum 15 ml per injection site in cattle over 150 kg, and 7 ml in pigs, calves, sheep and goats. Injection sites should be alternated.

- Cattle** 1 ml / 10 kg body mass (equivalent to 20 mg / kg body mass)
If necessary repeat after 72 hours.
- Sheep** 1 ml / 10 kg body mass (equivalent to 20 mg / kg body mass)
If necessary repeat after 60 hours.
- Goats** 1 ml / 10 kg body mass (equivalent to 20 mg / kg body mass)
If necessary repeat after 72 hours.
- Pigs** 1 ml / 10 kg body mass (equivalent to 20 mg / kg body mass)
If necessary repeat after 72 hours.

In case of tickborne gallsickness (anaplasmosis), repetitive treatments (10 - 14 day treatment period) are necessary to eliminate the parasite from carriers.

■ PRESENTATION

Amber coloured glass vials containing 100 ml, 250 ml and 500 ml sterile solution.

■ REGISTRATION HOLDER:

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000